Norton Audubon Hospital	Number Tested	Penicillins						Cephalosporins				Monobactam	Carbapenems		Aminoglycosides				Gram + Coverage					Others				
		Amoxicillin/Clavulanate	Ampicillin	Ampicillin/Sulbactam	Oxacilin	Penicilin	Piperacilin/Tazobactam	Cefazolin	Cefepime	Ceftazidime	Ceftriaxone	Aztreonam	Ertapenem	Meropenem	Amikacin	Gentamicin	Gentamicin Synergy	Trobramycin	Clindamycin [1, 2]	Erythromycin [2]	Vancomycin	Linezolid	Daptomycin	Ciprofloxacin	Levofloxacin	Nitrofurantoin [2]	Tetracycline	Trimeth/Sulfa
Acinetobacter baumannii complex	27	0	0	63	0	0		0	56	70		0	0	52	74	63		74	0	0	0	0	0					70
Other Acinetobacter species	11			91	0	0		0	91	91			0	91	91	82		82	0	0	0	0	0					82
Citrobacter freundii complex	43	0	0	0	0	0	93	0	95	67	67	71	100	100	100	91		91	0	0	0	0	0	86	88	93		81
Citrobacter koseri	30	97	0	93	0	0	100	90	100	100	97	100	100	100	100	100		100	0	0	0	0	0	70	70	85		77
Enterobacter cloacae complex	129	0	0	0	0	0	84	0	91	76	71	68	91	100	100	90		91	0	0	0	0	0	89	91	16		83
Escherichia coli	1282	80	45	52	0	0	96	58	90	89	87	87	99	100	99	90		89	0	0	0	0	0	70	71	96		70
Klebsiella aerogenes	48	0	0	0	0	0	88	0	98	81	73	80	98	100	100	98		98	0	0	0	0	0	96	96	9		96
Klebsiella oxytoca	60	83	0	70	0	0	93	10	95	93	88	89	98	96	100	93		93	0	0	0	0	0	92	98	92		92
Klebsiella pneumoniae	306	91	0	77	0	0	93	78	92	90	89	89	100	100	100	95		93	0	0	0	0	0	92	95	46		89
Morganella morganii	31	0	0	3	0	0	94	0	100	52	55	61	100	100	100	87		97	0	0	0	0	0	68	68	0		71
Proteus mirabilis	189	97	81	91	0	0	99	60	97	98	96	90	98	100	98	86		86	0	0	0	0	0	61	69	0	0	67
Proteus vulgaris	14	79	0	71	0	0	100	0	100	100	57	40	100	100	100	93		100	0	0	0	0	0	93	93	0	0	93
Providencia rettgeri	13	0	0	62	0	0	92	0	92	62	92	80	92	*	100	92		92	0	0	0	0	0	100	100	0	0	77
Providencia stuartii	16	0	0	12	0	0	100	0	100	94	88	100	94	*	100	0		0	0	0	0	0	0	50	56	0	0	88
Pseudomonas aeruginosa	248	0	0	0	0	0	86	0	85	84	0	74	0	89	98	85		96	0	0	0	0	0	82	79	0	0	0
Serratia marcescens	46	0	0	0	0	0	83	0	100	61	67	67	100	100	100	98		100	0	0	0	0	0	100	100	0		96
Stenotrophomonas maltophilia	24	0	0	0	0	0	0	0		38	0	0	0	0	0	0		0	0	0	0	0	0		79	0	0	96
Staphylococcus aureus	813	49			49			49				0							76	32	100	100	99				93	98
Methicillin-resistant S. aureus	428	0			0			0				0							69	11	100	100	99				95	96
Methicillin-susceptible S. aureus	412	100			100			100				0							83	53	100	100	100				92	100
Staphylococcus epidermidis	145	34			34			34				0							54	30	100	100	100			98	81	50
Staphylococcus haemolyticus	16	19			19			19				0							*	*	100	100	100			*	75	69
Staphylococcus lugdunensis	53	89			89			89				0							74	71	100	100	100			*	91	100
Staphylococcus simulans	11	82			82			82				0							*	*	100	100	100			*	91	100
Other coagulase-negative staphylococci	28	74			74			74				0							73	46	100	96	100			*	89	89
Enterococcus faecalis	148		99			99		0	0	0	0	0			0	0	72	0	0	26	91	100	100			*	22	0
Enterococcus faecium	26		27			23		0	0	0	0	0			0	0	81	0	0	4	35	100	88			*	27	0
Other Enterococcus species	12		75			67		0	0	0	0	0			0	0	100	0	0	20	75	100	100			*	17	0
Group B Streptococcus (S. agalactiae)	31		100			100					100	0							43	33	100				100		23	0
Streptococcus anginosus [3]	36		94			97					100	0							74	60	100				97		44	
Streptococcus constellatus [3]	30		87			93					100	0							80	69	100				97		70	
Streptococcus intermedius [3]	20		100			100					100	0							75	40	100				100		60	
Streptococcus pneumoniae	58	97				See Table II					See Table II	0							91	58	100				98		86	74
Streptococcus pyogenes	18		100			100					100	0							61	61	100				100		61	0
Viridans streptococci	35		74			71					97	0							80	51	100				91		43	
Aerococcus urinae	34		97			100					100	0							*	*	100				65		88	0

*Fewer than 10 isolates.

[1] MRSA: 7% inducible resistance, 24% constitutive resistance; MSSA: 13% inducible resistance, 4% constitutive resistance; Coag-neg Staph (all species): 6% inducible resistance, 32% constitutive resistance,
[2] Clindamycin and erythromycin data are based on non-urine isolates only. Nitrofurantoin susceptibility is based on urine isolates only.
[3] Together, the species S. anginosus, S. intermedius, and S. constellatus comprise the Streptococcus anginosus group.

Table II. Susceptibility rates of penicillin and ceftriaxone for *Streptococcus pneumoniae* (N = 58)	Penicillin - IV meningitis	Penicillin - IV non-meningitis	Penicillin - Oral	Ceftriaxone - IV meningitis	Ceftriaxone - IV non-meningitis
Percent Susceptible	73	100	73	93	100
Percent Intermediate	-	0	18	7	0
Percent Resistant	27	0	9	0	0