Norton Children's Hospital	Number Tested	Penicillins						Cephalosporins				Monobactam	Carbapenem	Aminoglycosides				Gram + Coverage					Others			
		Amoxicillin/Clavulanate	Ampicillin	Ampicillin/Sulbactam	Oxacillin	Penicillin	Piperacillin/Tazobactam	Cefazolin	Cefepime	Ceftazidime	Ceftriaxone	Aztreonam	Meropenem	Amikacin	Gentamicin	Gentamicin Synergy	Tobramycin	Clindamycin [1, 2]	Erythromycin [2]	Vancomycin	Linezolid	Daptomycin	Ciprofloxacin	Levofloxacin	Nitrofurantoin [2]	Trimeth/Sulfa
Enterobacter cloacae complex	46	0	0	0	0	0	74	0	96	67	59	65	100	100	98		98	0	0	0	0	0	100	100	22	91
Escherichia coli	501	83	41	45	0	0	97	57	95	95	94	93	100	99	90		91	0	0	0	0	0	87	87	98	69
Klebsiella oxytoca	26	85	0	58	0	0	92	12	100	100	92	88	100	100	100		100	0	0	0	0	0	96	96	60	100
Klebsiella pneumoniae	51	94	0	80	0	0	96	84	96	96	94	96	100	100	100		100	0	0	0	0	0	100	100	32	88
Proteus mirabilis	42	100	90	95	0	0	100	81	100	100	100	93	100	100	93		93	0	0	0	0	0	93	95	0	93
Pseudomonas aeruginosa	106	0	0	0	0	0	94	0	90	91	0	91	98	86	75		87	0	0	0	0	0	88	88	0	0
Serratia marcescens	21	0	0	0	0	0	71	0	100	48	52	50	100	100	90		81	0	0	0	0	0	86	95	0	90
Stenotrophomonas maltophilia	26	0	0	0	0	0	0	0		42	0	0	0	0	0		0	0	0	0	0	0		88	0	100
Staphylococcus aureus	508	51			51			51				0						79	39	100	100	100				99
Methicillin-resistant S. aureus	257	0			0			0				0						81	20	100	100	100				99
Methicillin-susceptible S. aureus	266	100			100			100				0						77	56	100	100	100				100
Staphylococcus epidermidis	71	42			42			42				0						54	30	100	97	100			100	69
Staphylococcus haemolyticus	13	31			31			31				0						*	*	100	100	100			91	77
Staphylococcus hominis	11	27			27			27				0						*	*	100	100	100			*	45
Staphylococcus simulans	16	69			69			69				0						*	*	100	100	100			100	100
Other coagulase-negative staphylococci	14	100			100			100				0						70	70	100	100	100			*	100
Enterococcus faecalis	89		100			100		0	0	0	0	0		0	0	83	0	0	26	100	100	100			98	0
Streptococcus anginosus [3]	17		100			100					94	0						82	71	100				100		
Streptococcus constellatus [3]	10		100			100					100	0						90	70	100				100		
Streptococcus intermedius [3]	14		86			86						0						93	79	100				100		
Streptococcus pneumoniae	53	91				See Table II					See Table II	0						86	47	100				100		57
Viridans streptococci	16		44			50					81	0						100	40	100				88		

*Fewer than 10 isolates.

[1] MRSA: 9% inducible resistance, 10% constitutive resistance; MSSA: 17% inducible resistance, 6% constitutive resistance; Coag-neg Staph (all species): 9% inducible resistance, 32% constitutive resistance.
[2] Clindamycin and erythromycin data are based on non-urine isolates only. Nitrofurantoin susceptibility is based on urine isolates only.
[3] Together, the species S. anginosus, S. intermedius, and S. constellatus comprise the Streptococcus anginosus group.

Table II. Susceptibility rates of penicillin and ceftriaxone for *Streptococcus pneumoniae* (N = 53)	Penicillin - IV meningitis	Penicillin - IV non-meningitis	Penicillin - Oral	Ceftriaxone - IV meningitis	Ceftriaxone - IV non-meningitis
Percent Susceptible	42	92	42	75	100
Percent Intermediate	-	6	30	25	0
Percent Resistant	58	2	28	0	0