

Norton Children's Medical Center	Number Tested	Penicillins						Cephalosporins				Monobactam	Carbapenem	Aminoglycosides				Gram + Coverage						Others			
		Amoxicillin/Clavulanate	Ampicillin	Ampicillin/Sulbactam	Penicillin	Oxacillin	Piperacillin/Tazobactam	Cefazolin	Cefepime	Ceftazidime	Ceftriaxone	Aztreonam	Meropenem	Amikacin	Gentamicin	Gentamicin Synergy	Tobramycin	Clindamycin [1, 2]	Erythromycin [2]	Vancomycin	Linezolid	Daptomycin	Ciprofloxacin	Levofloxacin	Nitrofurantoin [2]	Trimeth/Sulfa	
Escherichia coli	130	82	49	55	0	0	98	67	96	95	94	93	100	100	90	90	0	0	0	0	0	91	92	100	78		
Pseudomonas aeruginosa	11	0	0	0	0	0	91	0	100	100	0	*	100	100	91	100	0	0	0	0	0	100	100	0	0		
Staphylococcus aureus	63	59				59	59					0					89	42	100	100	100				98		
Methicillin-resistant S. aureus	24	0				0	0					0					90	10	100	100	100				96		
Methicillin-susceptible S. aureus	37	100				100	100					0					87	71	100	100	100				100		
Staphylococcus epidermidis	10	60				60	60					0					*	*	100	100	100			*	100		
Other coagulase-negative staphylococci	11	73				73	73					0					*	*	100	100	100			*	91		
Enterococcus faecalis	12		100		100		0	0	0	0	0	0	0	0	75	0	0	*	100	100	100			100	0		
*Fewer than 10 isolates.																											

[1] MRSA: 0% inducible resistance, 10% constitutive resistance; MSSA: 13% inducible resistance, 0% constitutive resistance.

[2] Clindamycin and erythromycin data are based on non-urine isolates only. Nitrofurantoin susceptibility is based on urine isolates only.