Outpatients	Number Tested	Penicillins						Cephalosporins				Monobactam	Carbapenems		Aminoglycosides				Gram + Coverage					Others				
		Amoxicillin/Clavulanate	Ampicillin	Ampicillin/Sulbactam	Oxacillin	Penicillin	Piperacillin/Tazobactam	Cefazolin	Cefepime	Ceftazidime	Ceftriaxone	Aztreonam	Ertapenem	Meropenem	Amikacin	Gentamicin	Gentamicin Synergy	Tobramycin	Clindamycin [1, 2]	Erythromycin [2]	Vancomycin	Linezolid	Daptomycin	Ciprofloxacin	Levofloxacin	Nitrofurantoin [2]	Tetracycline	Trimeth/Sulfa
Acinetobacter baumannii complex	28	0	0	86	0	0		0	89	89		0	0	92	96	93		93	0	0	0	0	0					82
Other Acinetobacter species	43	0	0	100	0	0		0	95	91		0	0	100	100	100		100	0	0	0	0	0					91
Aeromonas species	11									100	100			82										82				91
Citrobacter amalonaticus	14	71	14	50	0	0	100	14	100	100	71	83	100	*	100	100		100	0	0	0	0	0	93	93	18		100
Citrobacter freundii complex	128	0	0	0	0	0	97	0	100	88	84	86	98	100	100	99		98	0	0	0	0	0	97	98	92		88
Citrobacter koseri	142	97	0	95	0	0	99	95	99	98	99	98	100	100	100	99		99	0	0	0	0	0	99	99	86		99
Enterobacter cloacae complex	162	0	0	0	0	0	88	0	94	80	71	76	92	98	100	98		96	0	0	0	0	0	96	98	21		93
Escherichia coli	7994	86	55	58	0	0	98	68	96	96	95	95	99	100	99	93		94	0	0	0	0	0	86	86	99		79
Klebsiella aerogenes	167	0	0	0	0	0	95	0	99	89	85	92	99	100	100	98		98	0	0	0	0	0	98	99	19		98
Klebsiella oxytoca	155	94	0	71	0	0	97	20	99	99	97	95	100	100	100	99		96	0	0	0	0	0	94	95	85		90
Klebsiella pneumoniae	1135	96	0	87	0	0	97	89	98	98	98	98	99	97	99	99		98	0	0	0	0	0	98	99	38		92
Morganella morganii	44	0	0	7	0	0	100	0	100	75	77	66	100	100	100	100		100	0	0	0	0	0	95	96	0		89
Pantoea agglomerans group	11	91	0	100	0	0	100	0	100	100	100	86	100	100	100	100		100	0	0	0	0	0	100	100	*		100
Proteus mirabilis	503	99	90	97	0	0	100	77	99	99	99	96	100	100	99	97		97	0	0	0	0	0	93	95	0	0	87
Proteus vulgaris	10	80	0	70	0	0	100	0	100	90	10	*	100	100	100	100		100	0	0	0	0	0	100	100	0	0	90
Providencia rettgeri	13	0	0	46	0	0	100	0	100	31	100	67	100	*	100	100		100	0	0	0	0	0	92	92	0	0	92
Pseudomonas aeruginosa	292	0	0	0	0	0	97	0	93	96	0	91	0	94	94	83		94	0	0	0	0	0	91	92	0	0	0
Salmonella species	13		77		0	0					100								0	0	0	0	0	100				100
Serratia marcescens	42	0	0	0	0	0	69	0	100	69	67	68	100	100	100	100		93	0	0	0	0	0	90	98	0		100
Stenotrophomonas maltophilia	27	0	0	0	0	0	0	0		33	0	0	0	0	0	0		0	0	0	0	0	0		85			96
Staphylococcus aureus	1751	58			58			58				0							83	44	100	99	99				95	99
Methicillin-resistant S. aureus	735	0			0			0				0							83	17	100	99	99				96	99
Methicillin-susceptible S. aureus	1037	100			100			100				0							83	64	100	100	100				94	100
Staphylococcus capitis	13	92			92			92				0							*	*	100	100	100			*	100	85
Staphylococcus epidermidis	376	49			49			49				0							58	23	100	99	100			99	86	65
Staphylococcus haemolyticus	68	46			46			46				0							*	*	100	100	100			98	76	91
Staphylococcus hominis	27	59			59			59				0							*	*	100	100	100			100	89	89
Staphylococcus lugdunensis	160	95			95			95				0							88	84	100	100	100			100	94	99
Staphylococcus simulans	37	84			84			84				0							*	*	100	97	100			100	100	100
Other coagulase-negative staphylococci	25	72			72			72				0							*	*	100	100	100			100	96	92
Enterococcus faecalis	175		100			100		0	0	0	0	0			0	0	85	0	0	27	99	100	100			100	39	0
Other Enterococcus species	12		42			33		0	0	0	0	0			0	0	75	0	0	*	50	100	92			*	25	0
Group B Streptococcus (S. agalactiae)	525		100			100					100	0							49	33	100				99		15	0
Streptococcus anginosus [3]	64		98			98					100	0							86	56	100				100		30	
Streptococcus constellatus [3]	28		100			100					100	0							79	68	100				100		79	
Streptococcus intermedius [3]	27		100			100					100	0							81	67	100				100		59	
Streptococcus pneumoniae	75	99				See Table II					See Table II	0							90	44	100				100		92	72
Viridans streptococci	26		88			85					100	0							*	*	100				92		65	
Aerococcus urinae	71		96			96					92	0							*	*	100				63		90	0
*Fewer than 10 isolates.																												

[1] MRSA: 7% inducible resistance, 10% constitutive resistance; MSSA: 14% inducible resistance, 3% constitutive resistance; Coag-neg Staph (all species): 5% inducible resistance, 17% constitutive resistance.

[2] Clindamycin and erythromycin data are based on non-urine isolates only. Nitrofurantoin susceptibility is based on urine isolates only.

[3] Together, the species S. anginosus, S. intermedius, and S. constellatus comprise the Streptococcus anginosus group.

Table II. Susceptibility rates of penicillin and ceftriaxone for *Streptococcus pneumoniae* (N = 75)	Penicillin - IV meningitis	Penicillin - IV non-meningitis	Penicillin - Oral	Ceftriaxone - IV meningitis	Ceftriaxone - IV non-meningitis
Percent Susceptible	72	99	72	93	99
Percent Intermediate	-	0	14	6	1
Percent Resistant	28	1	14	1	0