Norton Hospital 2021	Number Tested	Penicillins						Cephalosporins					Monobactam	Carbapenems		Aminoglycosides				Gram + Coverage					Others				
		Amoxicillin/Clavulanate	Ampicillin	Ampicillin/Sulbactam	Oxacillin	Penicillin	Piperacillin/Tazobactam	Oral cephalosporins for uncomplicated UTI	Cefazolin	Cefepime	Ceftazidime	Ceftriaxone	Aztreonam	Ertapenem	Meropenem	Amikacin	Gentamicin	Gentamicin Synergy	Tobramycin	Clindamycin [1, 2]	Erythromycin [2]	Vancomycin	Linezolid	Daptomycin	Ciprofloxacin	Levofloxacin	Nitrofurantoin [2]	Tetracycline	Trimeth/Sulfa
Acinetobacter baumannii complex [3]	20	0	0	75	0	0			0	65	80		0	0	75	85	75		85	0	0	0	0	0	85	75			75
Citrobacter freundii complex [4]	29	0	0	0	0	0	93		0	97	83	79	83	97	97	100	90		93	0	0	0	0	0	90	93	89		83
Citrobacter koseri	19	89	0	84	0	0	95		89	100	100	89	95	100	100	100	100		100	0	0	0	0	0	89	100	62		95
Enterobacter cloacae complex [5]	89	0	0	0	0	0	79		0	88	71	56	65	82	97	100	99		98	0	0	0	0	0	97	100	21		94
Escherichia coli	965	85	47	55	0	0	97	86	66	92	91	87	88	99	99	99	91		91	0	0	0	0	0	74	75	97		72
Klebsiella aerogenes	41	0	0	0	0	0	88		0	98	85	80	83	95	100	100	100		100	0	0	0	0	0	95	95	18		98
Klebsiella oxytoca	53	98	0	81	0	0	100		15	98	94	96	94	100	100	100	100		96	0	0	0	0	0	98	98	80		96
Klebsiella pneumoniae	224	88	0	77	0	0	94	88	79	94	90	89	90	98	99	100	96		95	0	0	0	0	0	92	95	37		89
Morganella morganii	17	0	0	0	0	0	100		0	94	53	53	71	100	100	100	94		94	0	0	0	0	0	88	94	0		94
Proteus mirabilis	125	92	79	90	0	0	99	87	66	94	95	92	89	99	100	100	90		90	0	0	0	0	0	75	78	0	0	79
Proteus vulgaris	12	92	0	92	0	0	100		0	100	92	50	33	100	100	100	100		100	0	0	0	0	0	100	100	0	0	100
Pseudomonas aeruginosa	166	0	0	0	0	0	84		0	83	80	0	70	0	89	96	81		94	0	0	0	0	0	81	83	0	0	0
Serratia marcescens	47	0	0	0	0	0	57		0	98	43	53	38	100	100	98	96		91	0	0	0	0	0	100	100	0		98
Stenotrophomonas maltophilia	33	0	0	0	0	0	0		0		45	0	0	0	0	0	0		0	0	0	0	0	0		88		0	94
Staphylococcus aureus	682	47			47				47				0							77	33	100	100	99				91	98
Methicillin-resistant S. aureus	391	0			0				0				0							74	13	100	100	99				94	97
Methicillin-susceptible S. aureus	331	100			100				100				0							81	56	100	100	100				89	99
Staphylococcus capitis	10	100			100				100				0							*	*	100	100	100			*	100	100
Staphylococcus epidermidis	126	37			37				37				0							56	34	100	100	100			100	81	65
Staphylococcus haemolyticus	15	27			27				27				0							*	*	100	100	100			*	93	60
Staphylococcus lugdunensis	36	83			83				83				0							71	63	100	100	100			*	92	97
Other coagulase-negative staphylococci	25	50			50				50				0							46	40	100	100	100			*	80	84
Enterococcus faecalis	119		99			99			0	0	0	0	0			0	0	71	0	0	26	91	100	99			*	33	0
Enterococcus faecium	38		29			29			0	0	0	0	0			0	0	79	0	0	3	39	100	97			*	29	0
Other Enterococcus species	15		80			80			0	0	0	0	0			0	0	93	0	0	29	73	100	100			*	47	0
Group A Streptococcus (S. pyogenes)	11		100			100						100	0							55	45	100				100		36	0
Group B Streptococcus (S. agalactiae)	29		100			100						97	0							38	21	100				100		7	0
Streptococcus anginosus [6]	63		98			98						98	0							71	48	100				98		42	
Streptococcus constellatus [6]	37		97			100						100	0							65	51	100				100		59	
Streptococcus intermedius [6]	26		100			100						100	0							68	44	100				100		52	
Streptococcus pneumoniae	29	97				See Table II						See Table II	0							90	59	100				100		86	83
Viridans streptococci	46		59			54						91	0							83	24	100				93		61	
Aerococcus urinae	18		88			88						100	0							*	*	100				71		88	0

*Fewer than 10 isolates.

[1] MRSA: 5% inducible resistance, 21% constitutive resistance; MSSA: 12% inducible resistance, 7% constitutive resistance; Coag-neg Staph (all species): 4% inducible resistance, 36% constitutive resistance.
[2] Clindamycin and erythromycin data are based on non-urine isolates only. Nitrofurantoin susceptibility is based on urine isolates only.
[3] A. baumannii complex consists of the species A. baumannii, A. calcoaceticus, A. nosocomialis, and A. pitii.
[4] C. freundii complex consists of the species C. braakii, C. freundii, C. murliniae, C. sedlaki, C. werkmanii, and C. youngae.
[5] E. cloacae complex consists of the species E. asburiae, E. cloacae, E. hormaechei, E. kobei, E. ludwigii, and E. nimipressuralis.
[6] Together, the species S. anginosus, S. intermedius, and S. constellatus comprise the Streptococcus anginosus group.

Table II. Susceptibility rates of penicillin and ceftriaxone for *Streptococcus pneumoniae* (N = 29)	Penicillin - IV meningitis	Penicillin - IV non-meningitis	Penicillin - Oral	Ceftriaxone - IV meningitis	Ceftriaxone - IV non-meningitis
Percent Susceptible	83	97	83	93	97
Percent Intermediate	-	3	10	3	0
Percent Resistant	17	0	7	3	3